Trauma in Pregnancy

1-Nafiseh Bahrami, MD

Obstetrician-Gynecologist

2-Ali Ammini, MD

Obstetrician-Gynecologist, Associate Professor of Iran University of Medical Sciences,

Subspeciality of Prenatalogy, Head of Cynecology ward of Firouzgar Hospital

Abstract

Introduction: Trauma is leading complication and one of the causes of mortality of

pregnancy. Trauma can be lead to fetus death and threatening occurrences of

mother's life, whether resulting from vehicles accident or suicide and quarrel.

Discussion: It must be sured of opening of airway in pregnant women involved in

Trauma and necessity the complete oxygen should be prescribed or the patient is

under cannulation and mechanical treatment; since during pregnancy the womb is out

of pelvis and cause to pressure on the inferior vena cava and also pregnancy cause

to reduce of digestion system

movements and the pregnants's stomach must be assumed full, in cardiopulmonary

resuscitation and helping to protect of airway and service to vital organs the

mentioned items must be considered and in order to reject of any probable injury of

fetus and other processes of pregnancy, it is necessary that at least 4 hours of

monitoring to be done.

Conclusion: Surveys shows that the trauma resulting from pregnancy can be

preventable and with observance of safety measures and quick diagnosis of

suspecting cases of severe injury we able to reduce of mother and fetu's mortality.

Keyword: Pregnancy, blunt trauma, Penetrating trauma